Supplementary Table 1. #Enzian classification system (n=89).

	Overall (n =89)	EHSP (n=52)	EHDP1 (n=19)	EHDP2 (n=12)	EHDP3 (n=6)	*P* values
Peritoneum						
Absent	6 (6.7)	6 (11.5)	-	-	-	
P1	83 (93.3)	46 (88.5)	19 (100)	12 (100)	6 (100)	
P2	-	-	-	-	-	
P3	-	-	-	-	-	
Ovaries						
Absent	33 (37.1)	18 (34.6)	7 (36.8)	7 (58.3)	1 (16.7)	
O1	4 (4.5)	0	1 (5.3)	1 (8.3)	2 (33.3)	0.028*
O2	47 (52.8)	30 (57.7)	10 (52.6)	4 (33.3)	3 (50)	
O3	5 (5.6)	4 (7.7)	1 (5.3)	-	-	
Tubes						
Absent	-	-	-	-	-	
T1	4 (4.5)	4 (7.7)	-	-	-	0.050*
T2	64 (71.9)	40 (76.9)	12 (63.2)	10 (83.3)	2 (33.3)	
T3	21 (23.6)	8 (15.4)	7 (36.8)	2 (16.7)	4 (66.7)	
Compartments						
A (rectovaginal septum and vagina)						
Absent	67 (75.3)	43 (82.7)	11 (57.9)	10 (83.3)	3 (50)	
A1	4 (4.5)	1 (1.9)	2 (10.5)	0	1 (16.7)	
A2	17 (19.1)	8 (15.4)	5 (26.3)	2 (16.7)	2 (33.3)	0.223
A3	1 (1.1)	0	1 (5.3)	0	0	
B (uterosacral/cardinal ligaments, parametrium, pelvic sidewalls)						
Absent	-	-	-	-	-	
B1	1 (1.1)	1 (1.9)	0	0	0	
B2	88 (98.9)	51 (98.1)	19 (100)	12 (100)	6 (100)	
B3	-	-	-	-	-	
C (rectum)						
Absent	48 (53.9)	31 (59.6)	10 (52.6)	4 (33.3)	3 (50)	
C1	4 (4.5)	4 (7.7)	0	0	0	0.035*
C2	29 (32.6)	17 (32.7)	6 (31.6)	5 (41.7)	1 (16.7)	
C3	8 (9.0)	0	3 (15.8)	3 (25)	2 (33.3)	
Fa (adenomyosis)	89	52	19	12	6	0.044*
Fb (urinary bladder involvement)	3 (3.4)	1 (1.9)	2 (10.5)	0	0	0.266
Fi (other intestinal locations)	42 (47.2)	21 (40.4)	9 (47.4)	9 (75.0)	3 (50)	0.194
Fu (ureteric involvement with signs of obstruction)	9 (10.1)	2 (3.8)	3 (15.8)	0	4 (66.7)	0.001*
Fn (ipogastric nerve)	7 (7.9)	0	3 (15.8)	0	4 (66.7)	

Descriptive statistics are expressed as absolute and relative percentage frequencies. Statistically significant values were marked with an *.
EHSP: Extended hysterectomy with superficial parametrectomy, EHDP1: Extended hysterectomy with deep parametrectomy type 1, EHDP2: Extended hysterectomy with deep parametrectomy type 2, EHDP3: Extended hysterectomy with deep parametrectomy type 3.